

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 8 to nucleotide 2008;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone er311_20 deposited under accession number ATCC 98781;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone er311_20 deposited under accession number ATCC 98781;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:21.

2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.

3. A host cell transformed with the polynucleotide of claim 2.

4. The host cell of claim 3, wherein said cell is a mammalian cell.

5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:

- (a) growing a culture of a host cell transformed with the polynucleotide of claim 2 in a suitable culture medium; and
 - (b) purifying said protein from the culture.
- 6. A protein produced according to the process of claim 5.
- 7. An isolated polynucleotide encoding the protein of claim 6.
- 8. The polynucleotide of claim 7, wherein the polynucleotide comprises the cDNA insert of clone er311_20 deposited under accession number ATCC 98781.
- 9. A protein comprising an amino acid sequence selected from the group consisting of:
 - (a) the amino acid sequence of SEQ ID NO:22;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone er311_20 deposited under accession number ATCC 98781;the protein being substantially free from other mammalian proteins.
- 10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:22.
- 11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.
- 12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
 - (a) the nucleotide sequence of SEQ ID NO:65;
 - (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 66 to nucleotide 827;
 - (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 474 to nucleotide 827;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone d24_1 deposited under accession number ATCC 98817;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone d24_1 deposited under accession number ATCC 98817;
- (f) the nucleotide sequence of a mature protein coding sequence of clone d24_1 deposited under accession number ATCC 98817;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone d24_1 deposited under accession number ATCC 98817;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

13. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
 - (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
 - (c) the amino acid sequence encoded by the cDNA insert of clone d24_1 deposited under accession number ATCC 98817;
- the protein being substantially free from other mammalian proteins.